

# Anaphylaxis Emergency Action Plan

Patient Name: \_\_\_\_\_ Age: \_\_\_\_\_

Allergies: \_\_\_\_\_

Asthma  Yes (*high risk for severe reaction*)  No

Additional health problems besides anaphylaxis: \_\_\_\_\_

\_\_\_\_\_

Concurrent medications: \_\_\_\_\_

\_\_\_\_\_

### Symptoms of Anaphylaxis

MOUTH	itching, swelling of lips and/or tongue
THROAT*	itching, tightness/closure, hoarseness
SKIN	itching, hives, redness, swelling
GUT	vomiting, diarrhea, cramps
LUNGS*	shortness of breath, cough, wheeze
HEART*	weak pulse, dizziness, passing out

*Only a few symptoms may be present. Severity of symptoms can change quickly.  
\*Some symptoms can be life-threatening. ACT FAST!*

### Emergency Action Steps - DO NOT HESITATE TO GIVE EPINEPHRINE!

1. Inject epinephrine in thigh using (check one):

- 0.1 mg (16.5 lbs to less than 33 lbs) Specify brand: \_\_\_\_\_
- 0.15 mg (33 lbs to less than 66 lbs) Specify brand: \_\_\_\_\_
- 0.3 mg (66 lbs or more) Specify brand: \_\_\_\_\_

**IMPORTANT: ASTHMA INHALERS AND/OR ANTIHISTAMINES CAN'T BE DEPENDED ON IN ANAPHYLAXIS.**

2. Call 911 or emergency medical services (before calling contact)

3. Emergency contact #1: home \_\_\_\_\_ work \_\_\_\_\_ cell \_\_\_\_\_

Emergency contact #2: home \_\_\_\_\_ work \_\_\_\_\_ cell \_\_\_\_\_

Emergency contact #3: home \_\_\_\_\_ work \_\_\_\_\_ cell \_\_\_\_\_

Comments: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_  
Doctor's Signature/Date/Phone Number

\_\_\_\_\_  
Parent's Signature (for individuals under age 18 yrs)/Date